

# Molafos<sup>®</sup> Grazing Transition Pasture Supplement

Molafos Grazing Transition is a palatable supplement that is high in Molasses (Energy) with Magnesium, anionic salts and a sound mineral balance.

## Molafos Grazing Transition Uses

- ✓ For cattle close up to calving with good feed available, Molafos Grazing Transition should be fed for the last 2 weeks of pregnancy. Grazing Transition improves calcium metabolism, supplies magnesium and aids in reducing the incidence of many of the metabolic diseases that occur at or near calving.
- ✓ A Magnesium supplement in conditions where magnesium is either unavailable due to antagonists in the pasture or only available at low levels.
- ✓ Fed as a production supplement to Cattle, Sheep and Goats when there is a good body of forage available to grazing stock.
- ✓ Can be medicated with Monensin, Lasalocid, or Flavophospholipol to improve feed efficiency.

SUITABLE FOR SPECIES   

SUITABLE FOR CATEGORIES: Breeder, Backgrounder, Weaner, Finisher

## Molafos Grazing Transition Features

- Unique Molafos feed system. Excellent carrier of minerals, vitamins, supplements and medications.
- Inclusion of Magnesium and anionic salts, in a palatable carrier, ensures animals will accept these unpalatable but essential minerals.
- Ingredients are held in solution.
- Molasses based to supply sugars which stimulates appetite and improves urea utilisation.
- Flows at temperatures as low as -10°C.
- Contains a blend of Inorganic and Organic Minerals for improved availability and rumen bacterial health.
- Rain safe.

## Molafos Grazing Transition Benefits

- Assists when cattle on fresh green pasture are predisposed to a rapid fall in Magnesium levels, that can lead to nervous symptoms and even grass tetany.
- The anionic salts help improve blood Calcium and Magnesium levels by stimulating the animal to mobilise these minerals from their bones.
- Palatable and accepted by stock on all country types.
- Controllable intake and cost allows for a budget to be compiled for entire supplement program at commencement.
- Suitable for all types of cattle, sheep and goats regardless of body condition.
- Encourages stock to forage further and less selectively by promoting rumen bacteria function, ensuring enhanced forage utilisation, increased feed intake and digestion.
- Ultimate convenience with ease of use. Minimal labour required with flexibility in application.

## Directions for Use

- Pour into open troughs and allow cattle and sheep free access.
- Feed out 3 day's supply and do not refill until the 4th day.
- Once intake has stabilised feed out 7 day's supply and do not refill until the 8th day.
- Stock may need to be supplemented with salt (especially sheep) for 1 week to appease any salt craving.
- Do not feed to starving or thirsty stock.
- Place tubs close to areas where cattle will shelter during cold inclement weather to ensure they have access to the supplement at high risk times. Otherwise, locate feeding stations away from water to reduce intakes and encourage grazing out.

**Intakes – These products can be fed directly to livestock at recommended feeding rates.**

**Weaners: 0.5 to 1 kilogram / head / day**

**Adult Cattle: 1 to 1.5 kilograms / head / day**

**Sheep and Goats: 100 to 200 grams / head / day**

**1 litre Molafos = 1.32 kilograms**

### Molafos Grazing Transition – Analysis as Fed

Minimum Dry Matter	68%	Minimum Sulphur (S)	2%
Energy ME (Dry Matter)	9.5Mj/kg DM	Minimum Copper (Cu)	12mg/kg
Minimum Total Crude Protein	15.0%	Minimum Organic Copper (Cu)	6mg/kg
Minimum Crude Protein	3.0%	Minimum Cobalt (Co)	1.5mg/kg
Minimum Equiv Crude Protein	12%	Minimum Iodine (I)	10mg/kg
Maximum Urea	3.4%	Minimum Zinc (Zn)	300mg/kg
Minimum Calcium (Ca)	0.7%	Minimum Organic Zinc (Zn)	150mg/kg
Minimum Phosphorus(P)	0.5%	Minimum Manganese (Mn)	100mg/kg
Minimum Magnesium (Mg)	1.0%	Min Organic Manganese (Mn)	50mg/kg
Maximum Salt (NaCl)	3.0%	Minimum Selenium (Se)	2.2mg/kg

### Available Options – Monensin, Lasalocid and Flavophospholipol

#### WARNING

This product must be used in accordance with labeled directions. You must observe all directions, warnings and precautions noted in the instructions.

This product contains Urea. Urea can be toxic to Cattle and Sheep at levels as low as 0.25g of urea per kilogram of live weight. Livestock should be introduced to urea gradually. Avoid feeding urea to starving or thirsty livestock. Always ensure an adequate supply of forage and fresh water.

All medicated liquid supplements should be thoroughly agitated before use, either through mechanical mixing or recirculation on a daily basis.

Sodium Monensin is toxic to all Equines and Canines and should never be fed to these species.

Sodium Monensin and Lasalocid can be toxic to cattle and sheep and can be fatal if not fed in accordance with the instructions for use for this product.

**For further information please contact us on 1800 333 010 for customer service.**